

C^* -ALGEBRAS OF LABELLED GRAPHS

TERESA BATES & DAVID PASK

ABSTRACT. We describe a class of C^* -algebras which simultaneously generalise the ultragraph algebras of Tomforde and the shift space C^* -algebras of Matsumoto. In doing so we shed some new light on the different C^* -algebras that may be associated to a shift space. Finally, we show how to associate a simple C^* -algebra to an irreducible sofic shift.

KEYWORDS: C^* -algebras, labelled graph, ultragraph, Matsumoto algebra, shift space.

MSC (2000): 46L05, 37B10.

1. INTRODUCTION

The purpose of this paper is to introduce a class of C^* -algebras associated to labelled graphs. Our motivation is to provide a common framework for working with the ultragraph algebras of Tomforde (see [26, 27]) and the C^* -algebras associated to shift spaces studied by Matsumoto and Carlsen (see [14, 17, 6, 8] amongst others). Here a labelled graph (E, π) over an alphabet $\mathcal{A}$ is a directed graph E , together with a map $\pi : E^1 \rightarrow \mathcal{A}$. An ultragraph $\mathcal{G}$ is a particular example of a labelled graph (see Example 3.3 (ii)), and a shift space Λ has many presentations as a labelled graph (see [13], Example 3.3 (iii)). Hence it is natural to give our common framework in terms of labelled graphs.

To a two-sided shift space Λ over a finite alphabet, Matsumoto associates two C^* -algebras $\mathcal{O}_\Lambda$ and $\mathcal{O}_{\Lambda^*}$ generated by partial isometries (see [8]). Although $\mathcal{O}_\Lambda$ and $\mathcal{O}_{\Lambda^*}$ are generated by elements satisfying the same relations, it turns out that they are not isomorphic in general (see [8, Theorem 4.1]). This fact manifests itself in our realisation in section 6.2 of $\mathcal{O}_\Lambda$ and $\mathcal{O}_{\Lambda^*}$ as the C^* -algebras of the labelled graphs (E_Λ, π_Λ) and $(E_{\Lambda^*}, \pi_{\Lambda^*})$ respectively, which are not necessarily isomorphic as labelled graphs. Moreover, in Corollary 6.9 we show that using labelled graphs gives us the facility to canonically associate a simple C^* -algebra to an irreducible sofic shift (cf. [8, 6, 7]).

In fact we can associate a number of (possibly different) C^* -algebras to a labelled graph. This leads us to the notion of a labelled space, which we describe in section 3. Briefly, a labelled space $(E, \pi, \mathcal{C})$ consists of a labelled graph (E, π) together with a collection $\mathcal{C} \subseteq 2^{E^0}$ which plays the same role as $\mathcal{G}^0$ in [26] and is related to the abelian AF-subalgebra A_Λ (resp. A_{Λ^*}) in $\mathcal{O}_\Lambda$ (resp. $\mathcal{O}_{\Lambda^*}$) generated by the source projections.

In section 4 we define a representation of a labelled space in terms of partial isometries $\{s_a : a \in \mathcal{A}\}$ and projections $\{p_A : A \in \mathcal{C}\}$ subject to certain relations. Our relations generalise those found in [26, 14]. In order to build a nondegenerate C^* -algebra from a representation of $(E, \pi, \mathcal{C})$ it is necessary for $\mathcal{C}$ to be weakly left-resolving: a condition which is a generalisation of the left-resolving property for labelled graphs. Hence we may define $C^*(E, \pi, \mathcal{C})$ to be the C^* -algebra which is universal for representations of the weakly left-resolving labelled space $(E, \pi, \mathcal{C})$. Since any ultragraph has a natural realisation as a left-resolving labelled graph, the class of C^* -algebras of labelled spaces contains the ultragraph algebras (and hence, graph algebras and Exel-Laca algebras).

Date: March 17, 2005.

This research was supported by the University of Newcastle and the University of New South Wales.

In section 5 we give a version of the gauge invariant uniqueness theorem for $C^*(E, \pi, \mathcal{C})$ which will ultimately allow us to make the connection with the Matsumoto algebras.

In section 6 we give three applications of our uniqueness theorem: In section 6.1 we show how to construct a dual labelled space, which is the analogue of the higher block presentation of a shift space (cf. [13]). We give an isomorphism theorem for dual labelled spaces which is a generalisation of [2, Corollary 2.5] and forms a starting point for future work (see [4]). In section 6.2 we show that if $\mathcal{O}_\Lambda$ (resp. $\mathcal{O}_{\Lambda^*}$) has a gauge action, then it is isomorphic to the C^* -algebra of a certain labelled space. Then in section 6.3 we give necessary conditions for the C^* -algebra of a labelled space to be isomorphic to the C^* -algebra of the underlying directed graph. We then show how to associate a simple C^* -algebra to an irreducible shift space. By example, we show that in general the C^* -algebra of a labelled space will not be isomorphic to the C^* -algebra of any directed graph; hence labelled graph C^* -algebras form a strictly larger class of C^* -algebras than graph algebras.

Since we seek to generalise them, we begin by giving a brief description of Ultragraph algebras and Matsumoto algebras.

2. ULTRAGRAPH ALGEBRAS AND MATSUMOTO ALGEBRAS

2.1. Ultragraph Algebras. An ultragraph $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ consists of a countable set of vertices G^0 , a countable set of edges $\mathcal{G}^1$, and functions $s : \mathcal{G}^1 \rightarrow G^0$ and $r : \mathcal{G}^1 \rightarrow 2^{G^0}$. Let $\mathcal{G}^0$ be the smallest collection of 2^{G^0} which contains $s(e)$ and $r(e)$ for all $e \in \mathcal{G}^1$ and is closed under finite intersections and unions. The ultragraph algebra $C^*(\mathcal{G})$ is the universal C^* -algebra for Cuntz-Krieger $\mathcal{G}$ -families: collections of partial isometries $\{s_e : e \in \mathcal{G}^1\}$ with mutually orthogonal ranges, and projections $\{p_A : A \in \mathcal{G}^0\}$ satisfying the relations

1. $p_\emptyset = 0$, $p_A p_B = p_{A \cap B}$ and $p_{A \cup B} = p_A + p_B - p_{A \cap B}$ for all $A, B \in \mathcal{G}^0$
2. $s_e^* s_e = p_{r(e)}$ and $s_e s_e^* \leq p_{s(e)}$ for all $e \in \mathcal{G}^1$
3. $p_v = \sum_{s(e)=v} s_e s_e^*$ whenever $0 < |s^{-1}(v)| < \infty$

(see [26, Definition 2.7]). Recall that $v \in G^0$ is an infinite emitter if $|s^{-1}(v)| = \infty$. We say that $\mathcal{G}$ is *row-finite* if there are no infinite emitters and $r(e)$ is finite for all $e \in \mathcal{G}^1$.

Ultragraph algebras simultaneously generalise graph C^* -algebras and Exel-Laca algebras (see [26, Sections 3 and 4]). By [27, Corollary 5.5] there is a non row-finite ultragraph whose C^* -algebra is not Morita equivalent to a graph algebra or an Exel-Laca algebra.

2.2. Matsumoto Algebras. For an introduction to shift spaces we refer the reader to the excellent treatment in [13]. Let Λ be a two-sided shift space over a finite alphabet $\mathcal{A}$ and denote by Λ^* the collection of all finite blocks in Λ . Let

$$(1) \quad X_\Lambda = \{(x_i)_{i \geq 1} : (x_i)_{i \in \mathbf{Z}} \in \Lambda\}$$

denote the set of all right-infinite sequences in Λ .

Following [8] there are two C^* -algebras associated to Λ . Each C^* -algebra is generated by partial isometries $\{t_a : a \in \mathcal{A}\}$ subject to

$$(2) \quad \sum_{a \in \mathcal{A}} t_a t_a^* = 1, \text{ and } t_\alpha^* t_\alpha t_\beta = t_\beta t_{\alpha\beta}^* t_{\alpha\beta}, \text{ where } \alpha, \beta, \alpha\beta \in \Lambda^*.$$

As in [8] we denote by $\mathcal{O}_\Lambda$ the C^* -algebra defined directly on Hilbert space in [18, 19, 20] and by $\mathcal{O}_{\Lambda^*}$ the C^* -algebra defined using the Fock space construction in [14, 17, 16, 15, 21]. Because of the different ways in which the relations (2) are realised it turns out that $\mathcal{O}_\Lambda$ and $\mathcal{O}_{\Lambda^*}$ are not isomorphic in general (see [8, Section 6]).

There is a uniqueness theorem for $\mathcal{O}_\Lambda$ (resp. $\mathcal{O}_{\Lambda^*}$) when Λ satisfies condition (I) (resp. condition (I^*)) given in [8, Section 4] (resp. [8, Section 3]).

Condition (I): For $x \in X_\Lambda$ and $l \in \mathbf{N}$ put $\Lambda_l(x) = \{\mu \in \Lambda_l : \mu x \in X_\Lambda\}$. Two infinite paths $x, y \in X_\Lambda$ are *l-past equivalent* (written $x \sim_l y$) if $\Lambda_l(x) = \Lambda_l(y)$. The shift space X_Λ satisfies condition (I) if for any $l \in \mathbf{N}$ and $x \in X_\Lambda$ there exists $y \in X_\Lambda$ such that $y \neq x$, $y \sim_l x$.

Condition (I*): For $\omega \in \Lambda^*$ and $l \in \mathbf{N}$ we set $\Lambda_l(\omega) = \{\mu : |\mu| \leq l, \mu\omega \in \Lambda^*\}$. Two words $\mu, \nu \in \Lambda^*$ are said to be *l-past equivalent* (written $\mu \sim_l \nu$) if $\Lambda_l(\mu) = \Lambda_l(\nu)$. The subset $\Lambda_l^* \subseteq \Lambda^*$ is defined by

$$\Lambda_l^* := \{\omega \in \Lambda^* : |\{\mu \in \Lambda^* : \mu \sim_l \omega\}| < \infty\}.$$

The shift space Λ satisfies condition (I*) if for every $l \in \mathbf{N}$ and $\mu \in \Lambda_l^*$ there exist distinct words $\xi_1, \xi_2 \in \Lambda^*$ with $|\xi_1| = |\xi_2| = m$ such that

$$\mu \sim_l \xi_1 \gamma_1 \text{ and } \mu \sim_l \xi_2 \gamma_2$$

for some $\gamma_1, \gamma_2 \in \Lambda_{l+m}^*$.

Theorem 2.1. *Let Λ be a two-sided shift space over a finite alphabet which satisfies condition (I) (resp. (I*)). Then there is a strongly continuous action β (resp. β^*) of $\mathbf{T}$ on $\mathcal{O}_\Lambda$ (resp. $\mathcal{O}_{\Lambda^*}$) such that $\beta_z(t_a) = zt_a$ (resp. $\beta_z^*(t_a) = zt_a$) for all $a \in \mathcal{A}$ and $z \in \mathbf{T}$.*

Proof. That each β_z (resp. β_z^*) is an automorphism of $\mathcal{O}_\Lambda$ (resp. $\mathcal{O}_{\Lambda^*}$) for each $z \in \mathbf{T}$ follows from [8, Proposition 4.2] (resp. [8, Proposition 3.4]). A standard $\frac{\epsilon}{3}$ argument shows that β (resp. β^*) is strongly continuous. $\square$

In [21] Matsumoto defines λ -graph systems $\mathcal{L}_\Lambda$ and $\mathcal{L}_{\Lambda^*}$ associated to a two-sided shift space Λ together with corresponding C^* -algebras $\mathcal{O}_{\mathcal{L}_\Lambda}$ and $\mathcal{O}_{\mathcal{L}_{\Lambda^*}}$. By [8, Theorem 5.6] we see that if Λ satisfies condition (I) then $\mathcal{O}_\Lambda \cong \mathcal{O}_{\mathcal{L}_\Lambda}$ and if Λ satisfies condition (I*) then $\mathcal{O}_{\Lambda^*} \cong \mathcal{O}_{\mathcal{L}_{\Lambda^*}}$. Hence, for our purposes, it suffices to work with $\mathcal{O}_\Lambda$ and $\mathcal{O}_{\Lambda^*}$.

3. LABELLED SPACES

A directed graph E consists of a quadruple (E^0, E^1, r, s) where E^0 and E^1 are countable sets of vertices and edges respectively and $r, s : E^1 \rightarrow E^0$ are maps giving the direction of each edge. A path $\lambda = e_1 \dots e_n$ is a sequence of edges $e_i \in E^1$ such that $r(e_i) = s(e_{i+1})$ for $i = 1, \dots, n-1$. The collection of paths of length n in E is denoted E^n and the collection of all finite paths in E by E^* , so that $E^* = \cup_{n \geq 0} E^n$. The edge shift (X_E, σ_E) associated to a directed graph E with no sinks or sources is defined by:

$$X_E = \{x \in (E^1)^{\mathbf{Z}} : s(x_{i+1}) = r(x_i) \text{ for all } i \in \mathbf{Z}\} \text{ and } (\sigma_E x)_i = x_{i+1} \text{ for } i \in \mathbf{Z}.$$

The following definition is adapted from [13, Definition 3.1.1]:

Definition 3.1. A *labelled graph* (E, π) over an alphabet $\mathcal{A}$ consists of a directed graph E together with a labelling map $\pi : E^1 \rightarrow \mathcal{A}$.

Without loss of generality we may assume that the map π is onto. Given a labelled graph (E, π) such that every vertex in E emits and receives an edge, we may define a subshift $(X_{(E, \pi)}, \sigma)$ of $\mathcal{A}^{\mathbf{Z}}$ by

$$X_{(E, \pi)} = \{y \in \mathcal{A}^{\mathbf{Z}} : \text{there exists } x \in X_E \text{ such that } y_i = \pi(x_i) \text{ for all } i \in \mathbf{Z}\},$$

where σ is the shift map. The labelled graph (E, π) is said to be a *presentation* of the shift space $X = X_{(E, \pi)}$. As shown in [13, §3.1] a shift space may have many different presentations (see Examples 3.3 (ii), (vi), (vii)).

Let $\mathcal{A}^*$ be the collection of all *words* in the symbols of $\mathcal{A}$ (see [25, §0.2]). The map π extends naturally to a map $\pi : E^n \rightarrow \mathcal{A}^*$, where $n \geq 1$: for $\lambda = e_1 \dots e_n \in E^n$ put $\pi(\lambda) = \pi(e_1) \dots \pi(e_n)$; in this case the path $\lambda \in E^n$ is said to be a *representative* of the

labelled path $\pi(e_1) \dots \pi(e_n)$. Let $\mathcal{L}^n(E, \pi) = \pi(E^n)$ denote the collection of all labelled paths in (E, π) of length n , then $\mathcal{L}(E, \pi) = \cup_{n \geq 1} \mathcal{L}^n(E, \pi)$ denotes the collection of all words in the alphabet $\mathcal{A}$ which may be represented by paths in the labelled graph (E, π) . In this way π induces a map from the language $\cup_{n \geq 1} E^n$ of the subshift of finite type $\mathbf{X}_E$ associated to E into $\mathcal{L}(E, \pi)$, the language of the shift space $\mathbf{X}_{(E, \pi)}$ presented by (E, π) (see [13, §3]). The usual length function $|\cdot| : E^* \rightarrow \mathbf{N}$ transfers naturally over to $\mathcal{L}(E, \pi)$.

For α in $\mathcal{L}(E, \pi)$ we put

$$s_\pi(\alpha) = \{s(\lambda) \in E^0 : \pi(\lambda) = \alpha\} \text{ and } r_\pi(\alpha) = \{r(\lambda) \in E^0 : \pi(\lambda) = \alpha\},$$

so that $r_\pi, s_\pi : \mathcal{L}(E, \pi) \rightarrow 2^{E^0}$. We shall drop the subscript on r_π and s_π if the context in which it is being used is clear. For $\alpha, \beta \in \mathcal{L}(E, \pi)$ we have $\alpha\beta \in \mathcal{L}(E, \pi)$ if and only if $r(\alpha) \cap s(\beta) \neq \emptyset$.

Where possible we shall denote the elements of $\mathcal{A} = \mathcal{L}^1(E, \pi)$ as a, b , etc., elements of $\mathcal{L}(E, \pi)$ as α, β , etc., leaving e, f for elements of E^1 and λ, μ for elements of E^* .

Let (E, π) and (F, π') be graphs labelled by the same alphabet. A graph isomorphism $\phi : E \rightarrow F$ is a *labelled graph isomorphism* if $\pi'(\phi(e)) = \pi(e)$ for all $e \in E^1$ and we write $\phi : (E, \pi) \rightarrow (F, \pi')$.


Definition 3.2. The labelled graph (E, π) is *left-resolving* if for all $v \in E^0$ the map $\pi : r^{-1}(v) \rightarrow \mathcal{A}$ is injective.

The left-resolving condition ensures that for all $v \in E^0$ the labels $\{\pi(e) : r(e) = v\}$ of all incoming edges to v are all different. When (E, π) is left-resolving, if $\lambda, \mu \in \cup_{n \geq 1} E^n$ satisfy $\pi(\lambda) = \pi(\mu)$ and $r(\lambda) = r(\mu)$ then $\lambda = \mu$.

Examples 3.3. (i) Let E be a directed graph. Put $\mathcal{A} = E^1$ and let $\pi : E^1 \rightarrow E^1$ be the identity map (the *trivial labelling*) then (E, π) is a left-resolving labelled graph.

(ii) Let $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ be an ultragraph. Define $E = E_{\mathcal{G}}$ by putting $E^0 = G^0$, $E^1 = \{(s(e), w) : e \in \mathcal{G}^1, w \in r(e)\}$ and defining $r', s' : E^1 \rightarrow E^0$ by $s'(v, w) = v$, $r'(v, w) = w$. Set $\mathcal{A} = \mathcal{G}^1$ and define $\pi_{\mathcal{G}} : E^1 \rightarrow \mathcal{A}$ by $\pi(s(e), w) = e$. The resulting labelled graph $(E_{\mathcal{G}}, \pi_{\mathcal{G}})$ is left-resolving since the source map is single-valued. If $\mathcal{G}$ is row-finite then $E_{\mathcal{G}}$ is row-finite.

Conversely, given a left-resolving labelled graph (E, π) over an alphabet $\mathcal{A}$ where $s_\pi : \mathcal{L}(E, \pi) \rightarrow 2^{E^0}$ is single-valued, we can form a ultragraph $\mathcal{G}_E = (E^0, \mathcal{A}, r', s')$ with $s' = s_\pi$ and $r' = r_\pi$. If E is row finite then the ultragraph $\mathcal{G}_E$ is row-finite.



have the same language as the even shift Y since between any two 1's there must be an even number of 0's. Hence $\mathbf{X}_{(E_i, \pi_i)} = Y$ for $i = 1, 2, 3$ by [13, Proposition 1.3.4 (3)]. Only graphs (E_1, π_1) and (E_2, π_2) are left-resolving.

- (iv) Let E be a directed graph and Γ a group which acts on (the right of) E . Define $\pi_q : E^1 \rightarrow E^1/\Gamma$ by $\pi_q(e) = q(e)$ where $q : E^1 \rightarrow E^1/\Gamma$ is the quotient map. If the action of Γ is free on E^1 , then the resulting labelled graph (E, π_q) is left-resolving. More generally, if $p : F \rightarrow E$ is a graph morphism then there is a labelling $\pi_p : F^1 \rightarrow E^1$ given by $\pi_p(f) = p(f)$ for all $f \in F^1$. If p is a covering map then π_p is left-resolving.
- (v) Recall from [3, §3], that an out-splitting of a directed graph E is formed by a partition $\mathcal{P}$ of $s^{-1}(v)$ into $m(v) \geq 1$ non-empty subsets for each $v \in E^0$ (if $s^{-1}(v) = \emptyset$ then $m(v) = 0$). Given such a partition $\mathcal{P}$ one may construct a directed graph $E_s(\mathcal{P})$ where $E_s(\mathcal{P}^1) = \{e^j : e \in E^1, 1 \leq j \leq m(r(e))\} \cup \{e : m(r(e)) = 0\}$. Define $\pi : E_s(\mathcal{P})^1 \rightarrow E^1$ by $\pi(e^j) = e$ for $1 \leq j \leq m(r(e))$ and $\pi(e) = e$ if $m(r(e)) = 0$. For an in-splitting (see [3, §5]) of E using a partition $\mathcal{P}$, a similar construction also yields a labelled graph. However the resulting labelling π of the in-split graph $E_r(\mathcal{P})$ will not be left-resolving in general.
- (vi) Let Λ be a two-sided shift space over a finite alphabet $\mathcal{A}$ with X_Λ defined as in (1). Let $X_\Lambda^- = \{(x_i)_{i \leq 0} : (x_i)_{i \in \mathbb{Z}} \in \Lambda\}$ so that any element $x \in \Lambda$ may be written as $x = x^- x^+$. For arbitrary $x^+ \in X_\Lambda$ and $x^- \in X_\Lambda^-$ the bi-infinite sequence $y = x^- x^+$ may not belong to Λ . Define the past set of $t \in X_\Lambda$ as

$$P_\infty(t) = \{x^- \in X_\Lambda^- : x^- t \in \Lambda\}.$$

A shift is *sofic* if and only if the number of past sets is finite [11, 13].

For $s, t \in X_\Lambda$, we say that s is *past equivalent* to t (denoted $s \sim_\infty t$) if $P_\infty(s) = P_\infty(t)$. Evidently $X_\Lambda / \sim_\infty$ can be identified with Ω_Λ as described in [17, §2]. Define a labelled graph (E_Λ, π_Λ) as follows: let $E_\Lambda^0 = \{[v] : v \in X_\Lambda / \sim_\infty\}$, $E_\Lambda^1 = \{([v], a, [w]) : a \in \mathcal{A}, aw \sim_\infty v\}$ with $s([v], a, [w]) = [v]$ and $r([v], a, [w]) = [w]$. If $([v], a, [w]) \in E_\Lambda^1$ we put $\pi_\Lambda([v], a, [w]) = a$. The resulting left-resolving labelled graph is usually referred to as the *left-Krieger cover* of Λ and the construction is evidently independent of the choice of representatives (see [11]).

If Y is the even shift then (E_Y, π_Y) is labelled graph isomorphic to (E_2, π_2) in (iii) above. Let Z be shift over the alphabet $\{1, 2, 3, 4\}$ in which the words

$$\{12^k 1, 32^k 12, 32^k 13, 42^k 14 : k \geq 0\}$$

do not occur (see [8, §4]) then (E_Z, π_Z) has six vertices.

- (vii) Let Λ be a two-sided shift over a finite alphabet $\mathcal{A}$. We construct a variant of the predecessor graph $(E_{\Lambda^*}, \pi_{\Lambda^*})$ in the following way. For $\mu \in \Lambda^*$ we define

$$P(\mu) := \{\lambda : \lambda\mu \in \Lambda^*\}$$

and define an equivalence relation by $\mu \sim \nu$ if $P(\mu) = P(\nu)$. A shift is *sofic* if and only if the number of predecessor sets is finite [13].

Let Λ_∞^* denote those $\mu \in \Lambda^*$ which have an infinite equivalence class. Since $\mathcal{A}$ is finite $\Lambda_\infty^* / \sim$ can be identified with $\Omega_{\Lambda^*} = \lim_{\leftarrow} \Omega_t^*$ as described in [17, Section 2]. We set $E_{\Lambda^*}^0 = \Lambda_\infty^* / \sim$, $E_{\Lambda^*}^1 = \{([\mu], a, [\nu]) : a \in \mathcal{A}, [\mu] = [a\nu]\}$, $r([\mu], a, [\nu]) = [\nu]$ and $s([\mu], a, [\nu]) = [\mu]$. The labelling map is defined by $\pi_{\Lambda^*}([\mu], a, [\nu]) = a$. The resulting labelled graph is evidently left-resolving.

If Y is the even shift then (E_{Y^*}, π_{Y^*}) is labelled graph isomorphic to (E_2, π_2) in (iii) above (cf. [6, 17]). If Z is the sofic shift described in Example 3.3 (vi) then (E_{Z^*}, π_{Z^*}) has seven vertices and contains (E_Z, π_Z) as a subgraph.

Definition 3.4. Let (E, π) be a labelled graph. For $A \subseteq E^0$ and $\alpha \in \mathcal{L}(E, \pi)$ the *relative range of α with respect to A* is defined to be

$$r(A, \alpha) = \{r(\lambda) : \lambda \in E^*, \pi(\lambda) = \alpha, s(\lambda) \in A\}.$$

Remark 3.5. For any $A, B \subseteq E^0$ we have

$$r(A \cap B, \alpha) \subseteq r(A, \alpha) \cap r(B, \alpha) \text{ and } r(A \cup B, \alpha) = r(A, \alpha) \cup r(B, \alpha).$$

For all $A \subseteq E^0$ and $\alpha \in \mathcal{L}(E, \pi)$ we have $r(A, \alpha) = r(A \cap s(\alpha), \alpha)$.

A collection $\mathcal{C} \subseteq 2^{E^0}$ of subsets of E^0 is said to be *closed under relative ranges for (E, π)* if for all $A \in \mathcal{C}$ and $\alpha \in \mathcal{L}(E, \pi)$ we have $r(A, \alpha) \in \mathcal{C}$. If $\mathcal{C}$ is closed under relative ranges for (E, π) , contains $r(\alpha)$ for all $\alpha \in \mathcal{L}(E, \pi)$ and is also closed under finite intersections and unions, then we say that $\mathcal{C}$ is *accommodating* for (E, π) .

Definition 3.6. A *labelled space* consists of a triple $(E, \pi, \mathcal{C})$, where (E, π) is a labelled graph and $\mathcal{C}$ is accommodating for (E, π) .

Definition 3.7. A labelled space $(E, \pi, \mathcal{C})$ is *weakly left-resolving* if for every $A, B \in \mathcal{C}$ and every $\alpha \in \mathcal{L}(E, \pi)$ we have $r(A, \alpha) \cap r(B, \alpha) = r(A \cap B, \alpha)$.

In particular $(E, \pi, \mathcal{C})$ is weakly left-resolving if no pair of disjoint sets $A, B \in \mathcal{C}$ can emit paths λ, μ respectively with $\pi(\lambda) = \pi(\mu)$ and $r(\lambda) = r(\mu)$. If (E, π) is left-resolving then $(E, \pi, \mathcal{C})$ is weakly left-resolving for any $\mathcal{C}$. Evidently if $(E, \pi, \mathcal{C})$ is weakly left-resolving, then $(E, \pi, \mathcal{C}')$ is weakly left-resolving for any $\mathcal{C}' \subseteq \mathcal{C}$.

Consider the following subsets of 2^{E^0}

$$\begin{aligned} \mathcal{E} &= \{\{v\} : v \in E^0 \text{ is a source or a sink}\} \cup \{r(\alpha) : \alpha \in \mathcal{L}(E, \pi)\} \cup \{s(\alpha) : \alpha \in \mathcal{L}(E, \pi)\} \\ \mathcal{E}_- &= \{\{v\} : v \in E^0 \text{ is a sink}\} \cup \{r(\alpha) : \alpha \in \mathcal{L}(E, \pi)\}. \end{aligned}$$

The following definition is analogous to the definition of $\mathcal{G}^0$ in [26].

Definition 3.8. Let $\mathcal{E}^0$ (resp. $\mathcal{E}_-^0$) denote the smallest subset of 2^{E^0} containing $\mathcal{E}$ (resp. $\mathcal{E}_-$) which is accommodating for (E, π) .

Remark 3.9. For all $\alpha \in \mathcal{L}(E, \pi)$ we have $r(s(\alpha), \alpha) = r(\alpha)$; moreover if $(\alpha, \beta) \in \mathcal{L}^{(2)}(E, \pi)$ then $r(r(\alpha), \beta) = r(\alpha\beta)$. For $\alpha, \beta \in \mathcal{L}(E, \pi)$ with $\alpha\beta \in \mathcal{L}(E, \pi)$ and $A \subseteq E^0$ we have $r(r(A, \alpha), \beta) = r(A, \alpha\beta)$.

For labelled spaces $(E, \pi, \mathcal{E}^0)$ which are weakly left-resolving Remark 3.5 and Remark 3.9 show that to form $\mathcal{E}^0$ it suffices to form

$$\mathcal{E} \cup \{r(A, \alpha) : A \in \mathcal{E}, \alpha \in \mathcal{L}(E, \pi)\}$$

and then close under finite intersections and unions. To form $\mathcal{E}_-^0$, by Remark 3.5 it suffices to close $\mathcal{E}_-$ under finite intersections and unions. Evidently, $\mathcal{E}_-^0 \subseteq \mathcal{E}^0$; the containment can be strict, for instance this occurs when E has sources. One can show that $\mathcal{E}^0 = \mathcal{E}_-^0$ if and only if for every $\alpha \in \mathcal{L}(E, \pi)$, $s(\alpha)$ can be written as a finite union of sets of the form $\cap_{i=1}^n r(\beta_i)$. Since E^0 , $\mathcal{L}(E, \pi)$ and $\mathcal{E}$ are countable it follows that $\mathcal{E}^0$ and $\mathcal{E}_-^0$ are countable.

For $A \in 2^{E^0}$ and $n \geq 1$ let

$$L_A^n = \{\alpha \in \mathcal{L}^n(E, \pi) : A \cap s(\alpha) \neq \emptyset\}$$

denote those labelled paths of length n whose source intersects A nontrivially.

4. C*-ALGEBRAS OF LABELLED SPACES

Definition 4.1. Let $(E, \pi, \mathcal{C})$ be a weakly left-resolving labelled space. A representation of $(E, \pi, \mathcal{C})$ consists of projections $\{p_A : A \in \mathcal{C}\}$ and partial isometries $\{s_a : a \in \mathcal{L}^1(E, \pi)\}$ with the properties that

- (i) If $A, B \in \mathcal{C}$ then $p_A p_B = p_{A \cap B}$ and $p_{A \cup B} = p_A + p_B - p_{A \cap B}$, where $p_\emptyset = 0$.
- (ii) If $a \in \mathcal{L}^1(E, \pi)$ and $A \in \mathcal{C}$ then $p_A s_a = s_a p_{r(A, a)}$.
- (iii) If $a, b \in \mathcal{L}^1(E, \pi)$ then $s_a^* s_a = p_{r(a)}$ and $s_a^* s_b = 0$ unless $a = b$.
- (iv) For $A \in \mathcal{C}$, if L_A^1 is finite and non-empty we have

$$(3) \quad p_A = \sum_{a \in L_A^1} s_a p_{r(A, a)} s_a^*.$$

If $a, b \in \mathcal{L}^1(E, \pi)$ are such that $ab \in \mathcal{L}(E, \pi)$ then we have

$$(s_a^* s_a)(s_b s_b^*) = p_{r(a)} s_b s_b^* = s_b p_{r(r(a), b)} s_b^* = s_b s_b^* p_{r(a)} = (s_b s_b^*)(s_a^* s_a).$$

Hence $s_a s_b$ is a partial isometry which is nonzero if and only if s_a and s_b are. Therefore we may define $s_{ab} = s_a s_b$ and similarly define s_α for all $\alpha \in \mathcal{L}(E, \pi)$. One checks that Definition 4.1 (ii) holds for $\alpha \in \mathcal{L}(E, \pi)$, Definition 4.1 (iii) holds for $\alpha, \beta \in \mathcal{L}^n(E, \pi)$ for $n \geq 1$ and Definition 4.1 (iv) holds for $A \in \mathcal{C}$ with finite and nonempty L_A^n for $n \geq 1$. Then (cf. (2)) we have

$$(4) \quad s_\alpha^* s_\alpha s_\beta = p_{r(\alpha)} s_\beta = s_\beta p_{r(r(\alpha), \beta)} = s_\beta p_{r(\alpha \beta)} = s_\beta s_{\alpha \beta}^* s_{\alpha \beta}.$$

To justify the requirement that $(E, \pi, \mathcal{C})$ is weakly left-resolving in Definitions 4.1, consider the following: Let $\{p_A, s_a\}$ be a representation of $(E, \pi, \mathcal{C})$ in which $p_A \neq 0$ for all $A \in \mathcal{C}$. By Definition 4.1 (i) we have $(p_A - p_{A \cap B})(p_B - p_{A \cap B}) = 0$ for all $A, B \in \mathcal{C}$. Suppose, for contradiction, that there is $\alpha \in \mathcal{L}(E, \pi)$ such that $r(A, \alpha) \cap r(B, \alpha) \neq r(A \cap B, \alpha)$. From Definition 4.1 (iv) we have

$$p_A - p_{A \cap B} \geq s_\alpha (p_{r(A, \alpha)} - p_{r(A \cap B, \alpha)}) s_\alpha^* \text{ and } p_B - p_{A \cap B} \geq s_\alpha (p_{r(B, \alpha)} - p_{r(A \cap B, \alpha)}) s_\alpha^*$$

so $(p_A - p_{A \cap B})(p_B - p_{A \cap B}) \neq 0$, a contradiction. Thus a representation of $(E, \pi, \mathcal{C})$ will be degenerate if $(E, \pi, \mathcal{C})$ is not weakly left-resolving.

Relation (iv) in Definition 4.1 can make sense even if $A \in \mathcal{C}$ emits infinitely many edges in E : If there are only finitely many different labels attached to the edges which A emits then L_A^1 is finite. For directed graphs the analogue of equation (3) holds when a vertex has finite valency; when this is true at every vertex, the graph is called row-finite. With this in mind, we make the following definition:

Definition 4.2. Let $(E, \pi, \mathcal{C})$ be a labelled space. We say that $A \in \mathcal{C}$ is *singular* if L_A^1 is infinite. If no set $A \in \mathcal{C}$ is singular we say that $(E, \pi, \mathcal{C})$ is *set-finite*.

If $(E, \pi, \mathcal{C})$ is set-finite, then L_A^n is finite for all $A \in \mathcal{C}$ and all $n \geq 1$. In the examples below, the resulting labelled space will be set-finite whenever the original graph is row-finite.

Examples 4.3. (i) Let E be a directed graph with the trivial labelling π . Then $\mathcal{E}^0$ consists of all the finite subsets of E^0 . If E is row-finite then $(E, \pi, \mathcal{E}^0)$ and $(E, \pi, \mathcal{E}_-^0)$ are set-finite. One may show that a representation of $(E, \pi, \mathcal{E}^0)$ is a Cuntz-Krieger E -family and conversely (see [1, 2] for instance). If all sources in E have finite valency, then the $*$ -algebra generated by a representation of $(E, \pi, \mathcal{E}_-^0)$ contains a representation of $(E, \pi, \mathcal{E}^0)$. If there is a source $v \in E^0$ with infinite valency then there is no representative of p_v in the $*$ -algebra generated by a representation of $(E, \pi, \mathcal{E}_-^0)$.

- (ii) Under the identification of an ultragraph $\mathcal{G}$ with a labelled graph $(E_{\mathcal{G}}, \pi_{\mathcal{G}})$ we have $\mathcal{E}_{\mathcal{G}}^0 = \mathcal{G}^0$. Since $\mathcal{A} = \mathcal{G}^1$ a representation of $(E_{\mathcal{G}}, \pi_{\mathcal{G}}, \mathcal{E}_{\mathcal{G}}^0)$ is a Cuntz-Krieger $\mathcal{G}$ -family (see [26, Definition 2.7]). If $\mathcal{G}$ has sources which are singular then we get similar behaviour to that described in (i) above.
- (iii) In Examples 3.3 (iii) we have $\mathcal{E}_i^0, (\mathcal{E}_i^0)_- = 2^{E_i^0}$ for $i = 1, 2$ whereas $\{w\} \notin (\mathcal{E}_3^0)_-$. A representation of $(E_2, \pi_2, (\mathcal{E}_2^0)_-)$, is generated by partial isometries s_0, s_1 satisfying the relations in [14, Proposition 8.3] and [6, §2] for $\mathcal{O}_Y$, where Y is the even shift.
- (iv) A covering $p : F \rightarrow E$ of directed graphs yields a labelling $\pi_p : F^1 \rightarrow E^1$. We may identify $\mathcal{F}^0$ with the collection of inverse images of the finite subsets of E^0 . A representation of $(F, \pi_p, \mathcal{F}^0)$ is a Cuntz-Krieger E -family. If F has sources with infinite valency, then we get similar behaviour to that described in (i) above.
- (v) An outsplitting $E_s(\mathcal{P})$ of E gives rise to a labelling $\pi : E_s(\mathcal{P})^1 \rightarrow E^1$. If $\mathcal{P}$ is proper then we may identify $\mathcal{E}_s(\mathcal{P})^0$ with the collection of finite subsets of E^0 , and a representation of $(E_s(\mathcal{P}), \pi, \mathcal{E}_s(\mathcal{P})^0)$ is a Cuntz-Krieger E -family. If E has sources with infinite valency then, we get similar behaviour to that described in (i) above, even when the outsplitting is proper.
- (vi) An arbitrary shift $\Lambda \subseteq \mathcal{A}^{\mathbb{Z}}$ gives rise to a left-resolving labelled graph $(E_{\Lambda}, \pi_{\Lambda})$ with no sources or sinks. If $\mathcal{A}$ is finite then the generators of $\mathcal{O}_{\Lambda}$ form a representation of $(E_{\Lambda}, \pi_{\Lambda}, (\mathcal{E}_{\Lambda}^0)_-)$ (cf. [8, 14]).
- (vii) An arbitrary shift $\Lambda \subseteq \mathcal{A}^{\mathbb{Z}}$ gives rise to a left-resolving labelled graph $(E_{\Lambda^*}, \pi_{\Lambda^*})$ with no sources or sinks. If $\mathcal{A}$ is finite then the generators of $\mathcal{O}_{\Lambda^*}$ form a representation of $(E_{\Lambda^*}, \pi_{\Lambda^*}, (\mathcal{E}_{\Lambda^*}^0)_-)$ (cf. [8, 14]).

Examples 4.3 (i)-(v) show that it is possible for $\mathcal{E}^0$ and $\mathcal{E}_-^0$ to be different, but for the $*$ -algebras generated by representations of $(E, \pi, \mathcal{E}^0)$ and $(E, \pi, \mathcal{E}_-^0)$ to be the same.

Let $(E, \pi, \mathcal{C})$ be a labelled space. Let $\mathcal{C}^* = \mathcal{L}(E, \pi) \cup \mathcal{C}$ and extend r, s to $\mathcal{C}^*$ by $r(A) = A, s(A) = A$ for all $A \in \mathcal{C}$. For $A \in \mathcal{C}$, put $s_A = p_A$, so s_{β} is defined for all $\beta \in \mathcal{C}^*$.

Lemma 4.4. *Let $(E, \pi, \mathcal{C})$ be a weakly left-resolving labelled space and $\{s_a, p_A\}$ a representation of $(E, \pi, \mathcal{C})$. Then any nonzero product of s_a, p_A and s_b^* can be written as a finite combination of elements of the form $s_{\alpha} p_A s_{\beta}^*$ for some $A \in \mathcal{C}$, and $\alpha, \beta \in \mathcal{C}^*$ satisfying $A \subseteq r(\alpha) \cap r(\beta) \neq \emptyset$.*

Proof. Since $s_{\alpha} p_A s_{\beta}^* = s_{\alpha} p_{r(\alpha) \cap A \cap r(\beta)} s_{\beta}^*$ it follows that $s_{\alpha} p_A s_{\beta}^*$ is zero unless $A \cap r(\alpha) \cap r(\beta) \neq \emptyset$ and without loss of generality we may assume that $A \subseteq r(\alpha) \cap r(\beta)$. For $\alpha, \beta, \gamma, \delta \in \mathcal{L}(E, \pi)$ and $A, B \in \mathcal{C}$ we have

$$(5) \quad (s_{\alpha} p_A s_{\beta}^*) (s_{\gamma} p_B s_{\delta}^*) = \begin{cases} s_{\alpha \gamma'} p_{r(A, \gamma') \cap B} s_{\delta}^* & \text{if } \gamma = \beta \gamma' \\ s_{\alpha} p_{A \cap r(B, \beta')} s_{\delta \beta'}^* & \text{if } \beta = \gamma \beta' \\ s_{\alpha} p_{A \cap B} s_{\delta}^* & \text{if } \beta = \gamma \\ 0 & \text{otherwise} \end{cases}$$

To see this, suppose $\gamma = \beta \gamma'$ then as $A \subseteq r(\beta) \cap r(\alpha)$

$$\begin{aligned} s_{\alpha} p_A s_{\beta}^* s_{\gamma} p_B s_{\delta}^* &= s_{\alpha} p_A s_{\beta}^* s_{\beta \gamma'} p_B s_{\delta}^* = s_{\alpha} p_A p_{r(\beta)} s_{\gamma'} p_B s_{\delta}^* \\ &= s_{\alpha} p_A s_{\gamma'} p_B s_{\delta}^* = s_{\alpha \gamma'} p_{r(A, \gamma') \cap B} s_{\delta}^* \end{aligned}$$

A similar calculation gives the desired formulas in the cases $\beta = \gamma \beta'$ and $\beta = \gamma$. If β and γ have no common initial segment, then without loss of generality, assume that $\beta \in \mathcal{L}^n(E, \pi)$ and $\gamma \in \mathcal{L}^m(E, \pi)$ with $n > m$. Write $\beta = \beta' \beta''$ where $\beta' \in \mathcal{L}^m(E, \pi)$, and then by

Definition 4.1(iv) we have $s_\beta^* s_\gamma = s_{\beta'}^* s_{\beta'}^* s_\gamma = 0$ since $\beta' \neq \gamma$ and so $s_\alpha p_A s_\beta^* s_\gamma p_A s_\delta^* = 0$. By Definition 4.1 (i) and (ii) we may extend (5) to the case when $\alpha, \beta, \gamma, \delta \in \mathcal{C}^*$. $\square$

Theorem 4.5. *Let $(E, \pi, \mathcal{C})$ be a weakly left-resolving labelled space. There exists a C^* -algebra B generated by a universal representation of $\{s_a, p_A\}$ of $(E, \pi, \mathcal{C})$. Furthermore the s_a 's are nonzero and every p_A with $A \neq \emptyset$ is nonzero.*

Proof. Let $S_{(E, \pi, \mathcal{C})} := \{(\alpha, A, \beta) : \alpha, \beta \in \mathcal{C}^*, A \in \mathcal{C}, A \subseteq r(\alpha) \cap r(\beta)\}$ and let $k_{(E, \pi, \mathcal{C})}$ be the space of functions of finite support on $S_{(E, \pi, \mathcal{C})}$. The set of point masses $\{e_\tau : \tau \in S_{(E, \pi, \mathcal{C})}\}$ forms a basis for $k_{(E, \pi, \mathcal{C})}$. Set $(\alpha, A, \beta)^* := (\beta, A, \alpha)$; then thinking of $e_{(\alpha, A, \beta)}$ as $s_\alpha p_A s_\beta^*$ and using (5) we can define a multiplication with respect to which $k_{(E, \pi, \mathcal{C})}$ is a $*$ -algebra.

As a $*$ -algebra $k_{(E, \pi, \mathcal{C})}$ is generated by $q_A := e_{(A, A, A)}$ for $A \in \mathcal{C}$ and $t_a := e_{(a, r(a), r(a))}$ for $a \in \mathcal{L}^1(E, \pi)$. Our definition of multiplication ensures that properties (ii) and (iii) of Definition 4.1 hold; moreover $q_A q_B = q_{A \cap B}$. We mod out by the ideal J generated by the elements $q_{A \cup B} - q_A - q_B + q_{A \cap B}$ for $A, B \in \mathcal{C}$, and $q_A - \sum_{a \in L_A^1} s_a p_{r(A, a)} s_a^*$ for $A \in \mathcal{C}$ with L_A^1 nonempty and finite. Then the images r_A of q_A and u_a of t_a in $k_{(E, \pi, \mathcal{C})}/J$ form a representation of $(E, \pi, \mathcal{C})$ that generates $k_{(E, \pi, \mathcal{C})}/J$. The triple $(k_{(E, \pi, \mathcal{C})}/J, r_A, u_a)$ has the required universal property, but is not a C^* -algebra. Using a standard argument we can convert this triple to a C^* -algebra B satisfying the required properties (see [10, Theorem 2.1] for instance).

Now for each $a \in \mathcal{L}^1(E, \pi)$ and $e \in \pi^{-1}(a)$, let $\mathcal{H}_{(a, e)}$ be an infinite-dimensional Hilbert space. Also for each $v \in s(a)$ we define $\mathcal{H}_{(a, v)} := \bigoplus_{\{e: s(e)=v, \pi(e)=a\}} \mathcal{H}_{(a, e)}$. If v is a sink let $\mathcal{H}_v$ be an infinite-dimensional Hilbert space. For $A \in \mathcal{C}$ we define $\mathcal{H}_A := \bigoplus_{b \in L_A^1} \bigoplus_{v \in s(b) \cap A} \mathcal{H}_{(b, v)}$ and then note that each Hilbert space we have defined is a subspace of

$$\mathcal{H} := \left(\bigoplus_{a \in \mathcal{L}^1(E, \pi)} \bigoplus_{v \in s(a)} \mathcal{H}_{(a, v)} \right) \bigoplus_{\{v: s^{-1}(v)=\emptyset\}} \mathcal{H}_v.$$

For each $a \in \mathcal{L}^1(E, \pi)$, let S_a be a partial isometry with initial space $\mathcal{H}_{r(a)}$ and final space $\bigoplus_{v \in s(a)} \mathcal{H}_{(a, v)} \subseteq \mathcal{H}_{s(a)}$. For $A \in \mathcal{C}$, define P_A to be the projection of $\mathcal{H}$ onto $\mathcal{H}_A$, where this is interpreted as the zero projection when $A = \emptyset$.

It is easy to verify that since $(E, \pi, \mathcal{C})$ is weakly left-resolving, the operators $\{S_a, P_A\}$ form a representation of $(E, \pi, \mathcal{C})$ in which S_a, P_A are nonzero. By the universal property there exists a homomorphism $\pi_{S, P} : B \rightarrow C^*(\{S_a, P_A\})$. Since the S_a 's and P_A 's are nonzero, it follows that the s_a 's and p_A 's are also nonzero. $\square$

Definition 4.6. Let $(E, \pi, \mathcal{C})$ be a weakly left-resolving labelled space, then $C^*(E, \pi, \mathcal{C})$ is the universal C^* -algebra generated by a representation of $(E, \pi, \mathcal{C})$.

Let $(E, \pi, \mathcal{C})$ be a weakly left-resolving labelled space and $\{s_a, p_A\}$ be the universal representation of $(E, \pi, \mathcal{C})$, then by Lemma 4.4

$$\text{span} \{s_\alpha p_A s_\beta^* : \alpha, \beta \in \mathcal{L}(E, \pi), A \in \mathcal{C}, A \subseteq r(\alpha) \cap r(\beta)\}$$

is a dense $*$ -subalgebra of $C^*(E, \pi, \mathcal{C})$. The following result may be proved along the same lines as [26, Lemma 3.2].

Lemma 4.7. *Let $\mathcal{A}$ be finite, E have no sinks, and $(E, \pi, \mathcal{C})$ be a weakly left-resolving labelled space. Then $C^*(E, \pi, \mathcal{C})$ is unital.*

Proof. Observe that $\sum_{a \in \mathcal{A}} s_a s_a^*$ is a unit for $C^*(E, \pi, \mathcal{C})$. $\square$

Lemma 4.8. *If $\phi : (E, \pi) \rightarrow (F, \pi')$ is a labelled graph isomorphism, then for all $\mathcal{C}$ which are accommodating for (E, π) we have $C^*(E, \pi, \mathcal{C}) \cong C^*(F, \pi', \phi(\mathcal{C}))$.*

Proof. The map ϕ induces a bijection between the generators of $C^*(E, \pi, \mathcal{C})$ and $C^*(F, \pi', \phi(\mathcal{C}))$ and so by the universal property there are homomorphisms from one C^* -algebra to the other which are also inverses of each other. $\square$

5. GAUGE INVARIANT UNIQUENESS THEOREM

Let $\{s_a, p_A\}$ be the universal representation of $(E, \pi, \mathcal{C})$ which generates $C^*(E, \pi, \mathcal{C})$. For $z \in \mathbf{T}$, $a \in \mathcal{L}^1(E, \pi)$ and $A \in \mathcal{C}$ let

$$t_a := \gamma_z s_a = z s_a \text{ and } q_A := \gamma_z p_A = p_A$$

then the family $\{t_a, q_A\} \in C^*(E, \pi, \mathcal{C})$ is also a representation of $(E, \pi, \mathcal{C})$. By universality of $C^*(E, \pi, \mathcal{C})$ and a routine $\epsilon/3$ argument we see that γ extends to a strongly continuous action

$$\gamma : \mathbf{T} \rightarrow \text{Aut } C^*(E, \pi, \mathcal{C})$$

which we call the *gauge action*.

- Proposition 5.1.** i) Let E be a directed graph with the trivial labelling π . Then $C^*(E, \pi, \mathcal{E}^0) \cong C^*(E)$.
 ii) Let $\mathcal{G}$ be an ultragraph. Then $C^*(E_{\mathcal{G}}, \pi_{\mathcal{G}}, \mathcal{E}_{\mathcal{G}}^0) \cong C^*(\mathcal{G})$, where $(E_{\mathcal{G}}, \pi_{\mathcal{G}})$ is the labelled graph associated to $\mathcal{G}$.
 iii) Let $p : F \rightarrow E$ be a covering map with induced labelling $\pi_p : F^1 \rightarrow E^1$. Then $C^*(F, \pi_p, \mathcal{F}^0) \cong C^*(E)$.
 iv) Let E be a directed graph and let $E_s(\mathcal{P})$ be an outsplitting. Let π be the labelling of $E_s(\mathcal{P})$ induced by the outsplitting. If $\mathcal{P}$ is a proper partition then $C^*(E_s(\mathcal{P}), \pi, \mathcal{E}_s(\mathcal{P})^0) \cong C^*(E)$.

Proof. In each case the left hand side contains a generating set for the C^* -algebra on the right as shown in Examples 4.3. We apply the appropriate gauge-invariant uniqueness theorem for the algebra on the right hand side to obtain the isomorphism. $\square$

To establish connections with the Matsumoto algebras we need a version of the gauge-invariant uniqueness theorem for labelled graph algebras.

Lemma 5.2. Let $(E, \pi, \mathcal{C})$ be a weakly left-resolving labelled space, $\{s_a, p_A\}$ a representation of $(E, \pi, \mathcal{C})$, and $Y = \{s_{\alpha_i} p_{A_i} s_{\beta_i}^* : i = 1, \dots, N\}$ be a set of partial isometries in $C^*(E, \pi, \mathcal{C})$ which is closed under multiplication and taking adjoints. If q is a minimal projection in $C^*(Y)$ then either

- (i) $q = s_{\alpha_i} p_{A_i} s_{\alpha_i}^*$ for some $1 \leq i \leq N$
 (ii) $q = s_{\alpha_i} p_{A_i} s_{\alpha_i}^* - q'$ where $q' = \sum_{l=1}^m s_{\alpha_{k(l)}} p_{A_{k(l)}} s_{\alpha_{k(l)}}^*$ and $1 \leq i \leq N$; moreover there is a nonzero $r = s_{\alpha_i \beta} p_{r(A_i, \beta)} s_{\alpha_i \beta}^* \in C^*(E, \pi, \mathcal{C})$ such that $q'r = 0$ and $q \geq r$.

Proof. By equation (5) any projection in $C^*(Y)$ may be written as

$$\sum_{j=1}^n s_{\alpha_{i(j)}} p_{A_{i(j)}} s_{\alpha_{i(j)}}^* - \sum_{l=1}^m s_{\alpha_{k(l)}} p_{A_{k(l)}} s_{\alpha_{k(l)}}^*$$

where the projections in each sum are mutually orthogonal and for each l there is a unique j such that $s_{\alpha_{i(j)}} p_{A_{i(j)}} s_{\alpha_{i(j)}}^* \geq s_{\alpha_{k(l)}} p_{A_{k(l)}} s_{\alpha_{k(l)}}^*$.

If $q = \sum_{j=1}^n s_{\alpha_{i(j)}} p_{A_{i(j)}} s_{\alpha_{i(j)}}^* - \sum_{l=1}^m s_{\alpha_{k(l)}} p_{A_{k(l)}} s_{\alpha_{k(l)}}^*$ is a minimal projection in $C^*(Y)$ then we must have $n = 1$. If $m = 0$ then $q = s_{\alpha_i} p_{A_i} s_{\alpha_i}^*$ for some $1 \leq i \leq N$. If $m \neq 0$ then $q = s_{\alpha_i} p_{A_i} s_{\alpha_i}^* - q'$ where $q' = \sum_{l=1}^m s_{\alpha_{k(l)}} p_{A_{k(l)}} s_{\alpha_{k(l)}}^*$ and $1 \leq k \leq N$. Since q' is the sum of finitely many projections and $q \neq 0$ it follows by repeated use of Definition 4.1 (iv) that there is a nonzero $r = s_{\alpha_i \beta} p_{r(A_i, \beta)} s_{\alpha_i \beta}^*$ in $C^*(E, \pi, \mathcal{C})$ such that $r q' = 0$ and $q \geq r$. $\square$

Theorem 5.3. Let $(E, \pi, \mathcal{C})$ be a weakly left-resolving labelled space and let $\{S_a, P_A\}$ be a representation of $(E, \pi, \mathcal{C})$ on Hilbert space. Take $\pi_{S, P}$ to be the representation of $C^*(E, \pi, \mathcal{C})$ satisfying $\pi_{S, P}(s_a) = S_a$ and $\pi_{S, P}(p_A) = P_A$. Suppose that each P_A is non-zero

whenever $A \neq \emptyset$, and that there is a strongly continuous action β of $\mathbf{T}$ on $C^*(S_\alpha, P_A)$ such that for all $z \in \mathbf{T}$, $\beta_z \circ \pi_{S,P} = \pi_{S,P} \circ \gamma_z$. Then $\pi_{S,P}$ is faithful.

Proof. A straightforward argument along the lines of [22, Lemma 2.2.3] shows that

$$C^*(E, \pi, \mathcal{C})^\gamma = \overline{\text{span}}\{s_\alpha p_A s_\beta^* : \alpha, \beta \in \mathcal{L}^n(E, \pi) \text{ for some } n \text{ and } A \subseteq r(\alpha) \cap r(\beta)\}$$

where $C^*(E, \pi, \mathcal{C})^\gamma$ is the fixed point algebra of $C^*(E, \pi, \mathcal{C})$ under the gauge action γ . We claim that $C^*(E, \pi, \mathcal{C})^\gamma$ is AF. Let Y be a finite subset of $C^*(E, \pi, \mathcal{C})^\gamma$. Since $y \in Y$ may be approximated by a finite linear combination of elements of the form $s_\alpha p_A s_\beta^*$ where $|\alpha| = |\beta|$ we may assume that $Y = \{s_{\alpha_i} p_{A_i} s_{\beta_i}^* : |\alpha_i| = |\beta_i|, i = 1 \dots N\}$.

Let M be the length of the longest word in $\{\alpha_1, \dots, \alpha_N\}$. Let W denote the collection of all words in $\mathcal{L}(E, \pi)$ of length at most M that can be formed from composing subwords of $\alpha_1, \dots, \alpha_N, \beta_1, \dots, \beta_N$. Let $\mathcal{B}$ be the collection all finite intersections of $\{A_i\}_{i=1}^n$ and $\{r(A_i, \gamma) : 1 \leq i \leq N, \gamma \in W\}$. By equation (5) a non-zero product of elements of Y is of the form $s_\gamma p_A s_\delta^*$ where $\gamma, \delta \in W$ and $A \in \mathcal{B}$. Since W and $\mathcal{B}$ are finite it follows that $Y' = \{s_\gamma p_A s_\delta^* : \gamma, \delta \in W, A \in \mathcal{B}\}$ is finite, closed under adjoints and $C^*(Y) = C^*(Y')$. Hence we may assume that Y is closed under multiplication and taking adjoints. Thus $C^*(Y) = \overline{\text{span}}(Y)$ is finite dimensional and so $C^*(E, \pi, \mathcal{C})^\gamma$ is AF by [5, Theorem 2.2], establishing our claim.

To show that the canonical map $\pi_{S,P} : C^*(E, \pi, \mathcal{C}) \rightarrow C^*(S_\alpha, P_A)$ is injective on $C^*(E, \pi, \mathcal{C})^\gamma$ we write $C^*(E, \pi, \mathcal{C})^\gamma$ as $\bigcup C^*(Y_n)$ where $\{Y_n : n \geq 1\}$ is an increasing family of finite sets which are closed under multiplication and taking adjoints. Suppose, for contradiction, that $\pi_{S,P}$ is not faithful on $C^*(Y_n)$ for some n . Then its kernel is an ideal and so must contain a nonzero minimal projection q . If $Y_n = \{s_{\alpha_i} p_{A_i} s_{\beta_i}^* : i = 1 \dots, N(n)\}$ then by Lemma 5.2 either $q = s_{\alpha_i} p_{A_i} s_{\alpha_i}^*$ for some $1 \leq i \leq N(n)$ or $q = s_{\alpha_i} p_{A_i} s_{\alpha_i}^* - q'$ where $q' = \sum_{k=1}^m s_{\alpha_{i(k)}} p_{A_{i(k)}} s_{\alpha_{i(k)}}^*$ and $1 \leq i \leq N(n)$. In the first case $\pi_{S,P}(s_{\alpha_i} p_{A_i}) = S_{\alpha_i} P_{A_i}$ is a partial isometry with initial projection P_{A_i} and final projection $S_{\alpha_i} P_{A_i} S_{\alpha_i}^*$. But $P_{A_i} = \pi_{S,P}(p_{A_i}) \neq 0$ by hypothesis and so $\pi_{S,P}(q) = \pi_{S,P}(s_{\alpha_i} p_{A_i} s_{\alpha_i}^*) = S_{\alpha_i} P_{A_i} S_{\alpha_i}^* \neq 0$ which is a contradiction. In the second case by Lemma 5.2 (ii) there is $r = s_{\alpha_i \beta} p_{r(A_i, \beta)} s_{\alpha_i \beta}^*$ such that $q \geq r$ and $q'r = 0$. We may apply the above argument to show that $\pi_{S,P}(r) \neq 0$ and hence $\pi_{S,P}(q) \geq \pi_{S,P}(r) \neq 0$ which is also a contradiction. Hence $\pi_{S,P}$ is injective on $C^*(Y_n)$ and the result follows by arguments similar to those in [2, Theorem 2.1]. $\square$

6. APPLICATIONS

6.1. Dual Labelled Graphs. Let E have no sinks and (E, π) be a labelled graph over alphabet $\mathcal{A}$. From this data we may form the *dual labelled graph* $(\widehat{E}, \widehat{\pi})$ over alphabet $\widehat{\mathcal{A}} := \mathcal{L}^2(E, \pi)$ as follows: Let $\widehat{E}^0 = E^1$, $\widehat{E}^1 = E^2$ and the maps $r', s' : \widehat{E}^1 \rightarrow \widehat{E}^0$ be given by $r'(ef) = f$ and $s'(ef) = e$. The labelling $\widehat{\pi} : \widehat{E}^1 \rightarrow \widehat{\mathcal{A}}$ is induced by the original labelling, so that $\widehat{\pi}(ef) = \pi(e)\pi(f)$. For $ab \in \mathcal{L}^1(\widehat{E}, \widehat{\pi}) = \mathcal{L}^2(E, \pi)$ we have

$$r_{\widehat{\pi}}(ab) = \{f : \widehat{\pi}(ef) = ab\}, \text{ and } s_{\widehat{\pi}}(ab) = \{e : \widehat{\pi}(ef) = ab\}$$

and for $B \in 2^{E^1}$

$$r_{\widehat{\pi}}(B, ab) = \{f : \pi(ef) = ab, e \in B\}.$$

These maps extend naturally to $\mathcal{L}(\widehat{E}, \widehat{\pi}) = \bigcup_{n \geq 1} \mathcal{L}^n(\widehat{E}, \widehat{\pi})$ where for $n \geq 1$, $\mathcal{L}^n(\widehat{E}, \widehat{\pi})$ is identified with $\mathcal{L}^{n+1}(E, \pi)$. Consider the following subsets of 2^{E^1}

$$\begin{aligned} \widehat{\mathcal{E}} &= \{\{e\} : s(e) \text{ is a source}\} \cup \{r_{\widehat{\pi}}(\alpha) : \alpha \in \mathcal{L}(\widehat{E}, \widehat{\pi})\} \cup \{s_{\widehat{\pi}}(\alpha) : \alpha \in \mathcal{L}(\widehat{E}, \widehat{\pi})\} \\ \widehat{\mathcal{E}}_- &= \{r_{\widehat{\pi}}(\alpha) : \alpha \in \mathcal{L}(\widehat{E}, \widehat{\pi})\}. \end{aligned}$$

Let $\widehat{\mathcal{E}}^0$ (resp. $\widehat{\mathcal{E}}_-^0$) be the smallest collection of subsets of 2^{E^1} containing $\widehat{\mathcal{E}}$ (resp. $\widehat{\mathcal{E}}_-$) which is accommodating for $(\widehat{E}, \widehat{\pi})$. One checks easily that if $(E, \pi, \mathcal{C})$ is left-resolving, then $(\widehat{E}, \widehat{\pi}, \widehat{\mathcal{C}})$ is weakly left-resolving for $\mathcal{C} = \mathcal{E}^0, \mathcal{E}_-^0$.

For $B \in \widehat{\mathcal{E}}^0$ (resp. $B \in \widehat{\mathcal{E}}_-^0$) we set

$$\widehat{L}_B^1 = \{ab \in \mathcal{L}^1(\widehat{E}, \widehat{\pi}) : s_{\widehat{\pi}}(ab) \cap B \neq \emptyset\}.$$

If E has no sources and sinks, the shift $X_{(\widehat{E}, \widehat{\pi})}$ determined by the dual labelled graph $(\widehat{E}, \widehat{\pi})$ of (E, π) is the second higher block shift $X_{(E, \pi)}^{[2]}$ formed from $X_{(E, \pi)}$ (cf. [13, §1.4]).

Remarks 6.1. Suppose that $ab \in \mathcal{L}^2(E, \pi)$ then $c \in L_{r(ab)}^1$ if and only if $bc \in \widehat{L}_{r_{\widehat{\pi}}(ab)}^1$; moreover $r(r(ab), c) = r(s(r_{\widehat{\pi}}(ab)), bc)$. Suppose that $A \in \mathcal{E}^0$ (resp. $A \in \mathcal{E}_-^0$) then $a \in L_A^1$ and $ab \in \mathcal{L}^2(E, \pi)$ if and only if $ab \in \widehat{L}_{s^{-1}(A)}^1$.

Theorem 6.2. *Let (E, π) be a set-finite, left-resolving labelled graph with no sinks then $C^*(E, \pi, \mathcal{E}^0) \cong C^*(\widehat{E}, \widehat{\pi}, \widehat{\mathcal{E}}^0)$, moreover $C^*(E, \pi, \mathcal{E}_-^0) \cong C^*(\widehat{E}, \widehat{\pi}, \widehat{\mathcal{E}}_-^0)$.*

Proof. Let $\{s_a, p_A\}$ be a representation of $(E, \pi, \mathcal{E}^0)$ and $\{t_{ab}, q_B\}$ be a representation of $(\widehat{E}, \widehat{\pi}, \widehat{\mathcal{E}}^0)$. For $ab \in \mathcal{L}^1(\widehat{E}, \widehat{\pi})$ and $B \in \widehat{\mathcal{E}}^0$ let $T_{ab} = s_a s_b s_b^*$ and

$$Q_B := \sum_{ab \in \widehat{L}_B^1} s_{ab} p_{r(s(B), ab)} s_{ab}^*.$$

Since $(E, \pi, \mathcal{E}^0)$ is set-finite $(\widehat{E}, \widehat{\pi}, \widehat{\mathcal{E}}^0)$ is set-finite by Remarks 6.1 and so the above sum is finite. One checks that $\{T_{ab}, Q_B\}$ is a representation of $(\widehat{E}, \widehat{\pi}, \widehat{\mathcal{E}}^0)$.

By the universal property there is a homomorphism $\pi_{T, Q} : C^*(\widehat{E}, \widehat{\pi}, \widehat{\mathcal{E}}^0) \rightarrow C^*(E, \pi, \mathcal{E}^0)$ with $\pi_{T, Q}(t_{ab}) = T_{ab}$ and $\pi_{T, Q}(q_B) = Q_B$. Since $\pi_{T, Q}$ intertwines the respective gauge actions and $Q_B \neq 0$ it follows from Theorem 5.3 that $\pi_{T, Q}$ is faithful. We claim that $\pi_{T, Q}$ is surjective. For $a \in \mathcal{L}^1(E, \pi)$ we have

$$\begin{aligned} s_a &= s_a p_{r(a)} = s_a \sum_{b \in L_{r(a)}^1} s_b p_{r(r(a), b)} s_b^* = \sum_{b \in L_{r(a)}^1} s_a s_b s_b^* s_b p_{r(ab)} s_b^* \\ &= \sum_{b \in L_{r(a)}^1} s_a s_b s_b^* \sum_{c \in L_{r(ab)}^1} s_{bc} p_{r(r(ab), c)} s_{bc}^* \\ &= \sum_{b \in L_{r(a)}^1} T_{ab} \sum_{bc \in \widehat{L}_{r_{\widehat{\pi}}(ab)}^1} s_{bc} p_{r(s(r_{\widehat{\pi}}(ab)), bc)} s_{bc}^* \text{ by Remarks 6.1} \\ &= \sum_{b \in L_{r(a)}^1} T_{ab} Q_{r_{\widehat{\pi}}(ab)} \end{aligned}$$

and so $s_a \in C^*(T_{ab}, Q_B)$. For $A \in \mathcal{E}^0$, by Remarks 6.1 we have

$$\begin{aligned} p_A &= \sum_{a \in L_A^1} s_a p_{r(A, a)} s_a^* = \sum_{a \in L_A^1} s_a \sum_{b \in L_{r(A, a)}^1} s_b p_{r(r(A, a), b)} s_b^* s_a^* \\ &= \sum_{ab \in \widehat{L}_{s^{-1}(A)}^1} s_{ab} p_{r(A, ab)} s_{ab}^* = Q_{s^{-1}(A)} \end{aligned}$$

and so $p_A \in C^*(T_{ab}, Q_B)$ which establishes our claim. The second isomorphism is proved along similar lines. $\square$

6.2. Matsumoto Algebras.

Theorem 6.3. *Let Λ be a shift space over a finite alphabet $\mathcal{A}$ which satisfies condition (I) and has left-Krieger cover (E_Λ, π_Λ) then $\mathcal{O}_\Lambda \cong C^*(E_\Lambda, \pi_\Lambda, (\mathcal{E}_\Lambda)_-^0)$. Moreover, if Λ satisfies condition (I*), and has predecessor graph $(E_{\Lambda^*}, \pi_{\Lambda^*})$ then $\mathcal{O}_{\Lambda^*} \cong C^*(E_{\Lambda^*}, \pi_{\Lambda^*}, (\mathcal{E}_{\Lambda^*})_-^0)$.*

Proof. By definition every $A \in (\mathcal{E}_\Lambda)_-^0$ can be written as a union of sets of the form $A_j = \cap_{i=1}^{m(j)} r(\mu_i^j)$ for $j = 1, \dots, n$. For $\mu \in \Lambda^*$ let $q_{r(\mu)} = t_\mu^* t_\mu$, then since the projections $\{t_\mu^* t_\mu : \mu \in \Lambda^*\}$ are mutually commutative (see [17, p.686]) we may define $q_{r(\mu) \cap r(\nu)} = q_{r(\mu)} q_{r(\nu)}$, and hence define q_{A_j} for $1 \leq j \leq n$. By the inclusion-exclusion principle one may further define

$$q_A = \sum_{j=1}^n q_{A_j} - \sum_{j \neq k} q_{A_j} q_{A_k} + \dots + (-1)^{n+1} q_{A_1} \cdots q_{A_n}.$$

Using calculations along the lines of those in [14, §3] one checks that $\{t_a, q_A\}$ is a representation of $(E_\Lambda, \pi_\Lambda, (\mathcal{E}_\Lambda)_-^0)$.

Let $\{s_a, p_A\}$ be a representation of $(E_\Lambda, \pi_\Lambda, (\mathcal{E}_\Lambda)_-^0)$. By the universal property for $C^*(E_\Lambda, \pi_\Lambda, (\mathcal{E}_\Lambda)_-^0)$ there is a map $\psi_{t,q} : C^*(E_\Lambda, \pi_\Lambda, (\mathcal{E}_\Lambda)_-^0) \rightarrow \mathcal{O}_\Lambda$ such that $\psi_{t,q}(s_a) = t_a$ and $\psi_{t,q}(p_A) = q_A$, in particular $\psi_{t,q}$ is surjective.

Since Λ satisfies condition (I) it follows by Theorem 2.1 that $\mathcal{O}_\Lambda$ carries a strongly continuous action β of $\mathbf{T}$. Since $\beta_z \circ \psi_{t,q} = \psi_{t,q} \circ \gamma_z$ for all $z \in \mathbf{T}$ and $\psi_{t,q}(p_A) = q_A \neq 0$ it follows from Theorem 5.3 that $\psi_{t,q}$ is injective, which completes the proof of the first statement.

The second statement is proved similarly. $\square$

Remarks 6.4. (i) In [8, §5] a condition (*) is given under which for shift spaces Λ satisfying (*) conditions (I) and (I*) are equivalent and $\mathcal{O}_\Lambda \cong \mathcal{O}_{\Lambda^*}$. This suggests that if Λ satisfies (*) then (E_Λ, π_Λ) is labelled graph isomorphic to $(E_{\Lambda^*}, \pi_{\Lambda^*})$ and the isomorphism of $\mathcal{O}_\Lambda$ and $\mathcal{O}_{\Lambda^*}$ can be deduced from Theorem 4.8. However [8, Theorem 6.1] shows that, in general, $\mathcal{O}_\Lambda$ and $\mathcal{O}_{\Lambda^*}$ are not isomorphic. In particular, (E_Λ, π_Λ) and $(E_{\Lambda^*}, \pi_{\Lambda^*})$ are not labelled graph isomorphic in general.

(ii) The isomorphism of $C^*(E_\Lambda, \pi_\Lambda, (\mathcal{E}_\Lambda)_-^0)$ and $\mathcal{O}_\Lambda$ identifies $C^*(p_A : A \in (\mathcal{E}_\Lambda)_-^0)$ with $A_\Lambda \subset \mathcal{O}_\Lambda$. Recall from [17, Corollary 4.7] that $A_\Lambda \cong C(\Omega_\Lambda)$, hence we may think of the elements of $(\mathcal{E}_\Lambda)_-^0$ as indexing closed sets in Ω_Λ .

6.3. Finiteness Conditions.

Definition 6.5. A labelled graph (E, π) is *label-finite* if $|\pi^{-1}(a)| < \infty$ for all $a \in \mathcal{L}^1(E, \pi)$.

If (E, π) is label-finite then $\pi^{-1}(\alpha)$ is finite for all $\alpha \in \mathcal{L}(E, \pi)$ and so all sets in $\mathcal{E}^0$ are finite (and conversely). If (E, π) is label-finite then $(\widehat{E}, \widehat{\pi})$ is label-finite. If E is row-finite and (E, π) is label-finite then $(E, \pi, \mathcal{E}^0)$ is set-finite.

The following result generalises [2, Corollary 2.5] (see also [3, Remark 3.3 (i)]).

Theorem 6.6. *Let (E, π) be a row-finite left-resolving labelled graph which is label-finite and satisfies $\{v\} \in \mathcal{E}^0$ for all $v \in E^0$. Then $C^*(E, \pi, \mathcal{E}^0) \cong C^*(E)$; moreover if $\{v\} \in \mathcal{E}_-^0$ for all $v \in E^0$ then $C^*(E, \pi, \mathcal{E}_-^0) \cong C^*(E)$.*

Proof. Let $\{s_e, p_v\}$ be the canonical Cuntz-Krieger E -family and $\{t_a, q_A\}$ be the canonical generators of $C^*(E, \pi, \mathcal{E}^0)$. For $a \in \mathcal{L}^1(E, \pi)$ and $A \in \mathcal{E}^0$ let

$$T_a = \sum_{e \in E^1 : \pi(e) = a} s_e, \text{ and } Q_A = \sum_{v \in A} p_v.$$

The above sums make sense since (E, π) is label-finite. Since E is row-finite one may easily check that these operators define a representation of $(E, \pi, \mathcal{E}^0)$. By the universal property of $C^*(E, \pi, \mathcal{E}^0)$ there is a homomorphism $\psi_{T,Q} : C^*(E, \pi, \mathcal{E}^0) \rightarrow C^*(E)$ given by $\psi_{T,Q}(t_a) = T_a$ and $\psi_{T,Q}(q_A) = Q_A$ for all $a \in \mathcal{L}^1(E, \pi)$ and $A \in \mathcal{E}^0$.

Since $\{v\} \in \mathcal{E}^0$ for all $v \in E^0$, we have $p_v = Q_v \in C^*(T_a, Q_A)$ for all $v \in E^0$. Since our labelled graph is left-resolving we have $s_e = T_{\pi(e)}Q_{r(e)} \in C^*(T_a, Q_A)$ for all $e \in E^1$, and so $\psi_{T,Q}$ is surjective. The canonical gauge actions on $C^*(E)$ and $C^*(E, \pi, \mathcal{E}^0)$ satisfy the required properties and $\psi_{T,Q}(q_A) = Q_A \neq 0$ for all $A \in \mathcal{E}^0$, so $\psi_{T,Q}$ is an isomorphism by Theorem 5.3.

The proof of the second isomorphism is essentially the same. $\square$

Corollary 6.7. *Let $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ be a row-finite ultragraph then $C^*(\mathcal{G}) \cong C^*(E_{\mathcal{G}})$ where $E_{\mathcal{G}}$ is the underlying directed graph of $\mathcal{G}$.*

Proof. From Examples 3.3 (ii) a row-finite ultragraph $\mathcal{G}$ may be realised as a row-finite left-resolving labelled graph $(E_{\mathcal{G}}, \pi_{\mathcal{G}})$. As $E_{\mathcal{G}}$ is row-finite it follows that $(E_{\mathcal{G}}, \pi_{\mathcal{G}})$ is label-finite. Since the source map is single-valued it follows that $v \in \mathcal{E}_{\mathcal{G}}^0$ for all $v \in G^0 = E_{\mathcal{G}}^0$ and hence the result follows from Theorem 6.6. $\square$

The following result was first observed in [6, Theorem 3.5] (see also [24, Theorem 4.4.4]).

Corollary 6.8. *Let Λ be a sofic shift over a finite alphabet then*

$$\mathcal{O}_{\Lambda} \cong C^*(E_{\Lambda})$$

where $(E_{\Lambda}, \pi_{\Lambda})$ is the left-Krieger cover of Λ .

Proof. As E_{Λ}^0 is finite and each $v \in E_{\Lambda}^0$ has a different past there are $\alpha_v \in \mathcal{L}(E_{\Lambda}, \pi_{\Lambda})$ with $r_{\pi_{\Lambda}}(\alpha_v) = \{v\}$. Hence $\{v\} \in (\mathcal{E}_{\Lambda}^0)_{-}$ for all $v \in E_{\Lambda}^0$. The result follows by Theorem 6.6. $\square$

From [13, Theorem 3.3.18] any two minimal left-resolving representations (E, π) , (F, π') of an irreducible sofic shift are labelled graph isomorphic and so $C^*(E, \pi, \mathcal{E}_{-}^0) \cong C^*(F, \pi', \mathcal{F}_{-}^0)$ by Lemma 4.8. Moreover, one may use the minimality of the representation to show that the underlying graph E is irreducible (cf. [13, Lemma 3.3.10]). Hence we have:

Corollary 6.9. *Let (E, π) be a minimal left-resolving presentation of an irreducible sofic shift over a finite alphabet, then $C^*(E, \pi, \mathcal{E}_{-}^0) \cong C^*(E, \pi, \mathcal{E}^0)$ is simple.*

Remark 6.10. Recall that the graph (E_2, π_2) in Examples 3.3 (ii) is the left-Krieger cover of the even shift Y . Although Y is irreducible, (E_2, π_2) is not a minimal left-resolving presentation of Y and $\mathcal{O}_Y \cong C^*(E_2)$ is not simple. However the graph (E_1, π_1) Examples 3.3 (ii) is a minimal left-resolving cover of Y and so

$$C^*(E_1, \pi_1, \mathcal{E}_{-}^0) \cong C^*(E_1, \pi_1, \mathcal{E}^0) \cong C^*(E_1)$$

is simple. Similarly $C^*(E_Z, \pi_Z, \mathcal{E}_{-}^0) \cong C^*(E_Z)$ is simple where Z is the irreducible shift introduced in Examples 3.3 (vi).

Thus, if one wishes to associate a simple C^* -algebra to an irreducible sofic shift Λ , then one should use the minimal left-resolving presentation of Λ . (cf. [6, 7]).

For a general shift space Λ , either $(E_{\Lambda}, \pi_{\Lambda})$ will not be row-finite or there will be $v \in E_{\Lambda}^0$ with $v \notin (\mathcal{E}_{\Lambda}^0)_{-}$. This indicates that the C^* -algebras corresponding to presentations of such shift spaces will not be Morita equivalent to graph algebras. The shift associated to a certain Shannon graph (see [20, Theorem 7.7]) provides such an example.

REFERENCES

- [1] T. Bates, J.-H. Hong, I. Raeburn and W. Szymański. *The ideal structure of the C^* -algebras of infinite graphs*. Illinois J. Math **46** (2002), 1159–1176.
- [2] T. Bates, D. Pask, I. Raeburn and W. Szymański. *The C^* -algebras of row-finite graphs*. New York J. Math. **6** (2000), 307–324.
- [3] T. Bates and D. Pask. *Flow equivalence of graph algebras*. Ergod. Th. & Dynam. Sys., **24** (2004), 367–382.
- [4] T. Bates and D. Pask. *Flow equivalence of labelled graph algebras*. In preparation.
- [5] O. Bratteli. *Inductive limits of finite dimensional C^* -algebras*. Trans. Amer. Math. Soc., **171** (1972), 195–234.
- [6] T. Carlsen. *On C^* -algebras Associated with Sofic Shifts*. J. Operator Theory **49** (2003), 203–212.
- [7] T. Carlsen. *Symbolic dynamics, partial dynamical systems, boolean algebras and C^* -algebras generated by partial isometries*, preprint Univ. Oslo, (2004).
- [8] T. Carlsen and K. Matsumoto. *Some remarks on the C^* -algebras associated with subshifts*. Math. Scand. **95** (2004), 145–160.
- [9] K. Deicke, D. Pask and I. Raeburn, *Coverings of directed graphs and crossed products of C^* -algebras by coactions of homogeneous spaces*. Internat. J. Math., **14** (2003), 773–789.
- [10] A. an Huef and I. Raeburn. *The ideal structure of Cuntz-Krieger algebras*. Ergod. Th. & Dynam. Sys. **17** (1997), 611–624.
- [11] W. Krieger. *Sofic Systems I*. Israel J. Math., **48**, (1984), 305–330.
- [12] A. Kumjian, D. Pask and I. Raeburn. *Cuntz-Krieger algebras of directed graphs*. Pacific J. Math. **184** (1998), 161–174.
- [13] D. Lind and B. Marcus. *An Introduction to Symbolic Dynamics and Coding*, CUP, 1995.
- [14] K. Matsumoto. *On C^* -algebras associated with subshifts*. Internat. J. Math. **8**, (1997), 357–374.
- [15] K. Matsumoto. *K -theory for C^* -algebras associated with subshifts*. Math. Scand. **82**, (1998), 237–255.
- [16] K. Matsumoto. *Relations among generators of C^* -algebras associated with subshifts*, Internat. J. Math. **10** (1999), 385–405.
- [17] K. Matsumoto. *Dimension groups for subshifts and simplicity of the associated C^* -algebras*. J. Math. Soc. Japan **51** (1999), 679–697.
- [18] K. Matsumoto. *On automorphisms of C^* -algebras associated with subshifts*. J. Operator Theory **44**, (2000), 91–112.
- [19] K. Matsumoto. *Bowen-Franks groups for subshifts and Ext-groups for C^* -algebras*. K-Theory **23**, (2001), 67–104.
- [20] K. Matsumoto. *Stabilized C^* -algebras associated with presentations of subshifts*. Documenta Math. **7**, (2002), 1–30.
- [21] K. Matsumoto. *Stabilized C^* -algebras constructed from symbolic dynamical systems*. Ergodic Theory Dynam. Sys. **20**, (2000), 821–841.
- [22] D. Pask and I. Raeburn, *On the K -theory of Cuntz-Krieger algebras*. Publ. RIMS Kyoto Univ. **32** (1996), 415–443.
- [23] I. Raeburn and W. Szymański. *Cuntz-Krieger algebras of infinite graphs and matrices*. Trans. AMS, **356**, (2004) 39–59.
- [24] J. Samuel. *Graphs, Sofic Shifts and C^* -algebras I*. Preprint Univ. Victoria, (1998).
- [25] R.G. Taylor. *Models of computation and formal languages*, Oxford University Press, 1998.
- [26] M. Tomforde. *A unified approach to Exel-Laca algebras and C^* -algebras associated to graphs*. J. Operator Theory **50** (2003), 345–368.
- [27] M. Tomforde. *Simplicity of ultragraph algebras*. Indiana Univ. Math. J. **52** (2003), 901–926.

SCHOOL OF MATHEMATICS, UNSW, SYDNEY 2052, AUSTRALIA
E-mail address: teresa@unsw.edu.au

MATHEMATICS, UNIVERSITY OF NEWCASTLE, NSW 2308, AUSTRALIA
E-mail address: david.pask@newcastle.edu.au